Synthesis and reactivity of dichloroboryl complexes of platinum(II)

Jonathan P. H. Charmant, Cheng Fan, Nicholas C. Norman* and Paul G. Pringle*

Received 4th September 2006, Accepted 16th October 2006 First published as an Advance Article on the web 6th November 2006 DOI: 10.1039/b612039n

The reaction between $[Pt(nbe)_3]$ (nbe = norbornene), two equivalents of the phosphines PPh₃, PMePh₂ or PMe₂Ph and 1 equivalent of BCl₃ affords the platinum dichloroboryl species [PtCl(BCl₂)(PPh₃)₂], [PtCl(BCl₂)(PMePh₂)₂] and [PtCl(BCl₂)(PMe₂Ph)₂]. All three complexes were characterised by X-ray crystallography and reveal that the boryl group lies trans to the chloride. With PMe₃ as the phosphine, the complex $[PtCl(BCl_2)(PMe_3)_2]$ is isolated in high yield as a white crystalline powder although crystals suitable for X-ray crystallography were not obtained. Crystals were obtained of a product shown by X-ray crystallography to be the unusual dinuclear species $[Pt_2(BCl_2)_2(PMe_3)_4(\mu-Cl)][BCl_4]$ which reveals an arrangement in which two square planar platinum(II) centres are linked by a single bridging chloride which is *trans* to a BCl₂ group on each platinum centre. The reaction of $[PtCl(BCl_2)(PMe_3)_2]$ with NEt₃ or pyridine (py) affords the adducts [PtCl{BCl₂(NEt₃)}(PMe₃)₂] and [PtCl{BCl₂(py)}(PMe₃)₂], respectively, both characterised spectroscopically. The reaction between $[PtCl(BCl_2)(PMe_3)_2]$ and either 4 equivalents of NHEt₂ or piperidine (pipH) results in the mono-substituted boryl species $[PtCl{BCl(NEt_2)}(PMe_3)_2]$ and $[PtCl{BCl(pip)}(PMe_3)_2]$, respectively, the former characterised by X-ray crystallography. Treatment of either [PtCl(BCl₂)(PMe₃)₂] (in the presence of excess NEt₃) or $[PtCl{BCl(NEt_2)}(PMe_3)_2]$ with catechol affords the B(cat) (cat = catecholate) derivative $[PtCl{B(cat)}(PMe_3)_2]$ which is also formed in the reaction between $[Pt(PMe_3)_4]$ and ClB(cat) and also from the slow decomposition of $[Pt{B(cat)}_2(PMe_3)_2]$ in dichloromethane over a period of months. The compound $[Pt{B(cat)}_2(PMe_3)_2]$ was prepared from the reaction between $[Pt(PMe_3)_4]$ and $B_2(cat)_2$.

Introduction

Transition metal boryl compounds¹ remain a focus of continued interest not least because of their involvement in various metal catalysed diboration and other borylation reactions.² In most examples, the boron of the boryl ligand is bonded to a single metal centre and two additional atoms which are usually good π -donors such as oxygen or nitrogen incorporated into diolato or amido groups respectively; the B(cat) group, where cat = catecholate, is one of the most studied.¹ More recent investigations, notably by Braunschweig *et al.* and Aldridge *et al.*, have focussed on more reactive boryl species in which the R groups of the BR₂ ligand are hydrocarbyl and/or halogeno groups.

With particular reference to dihaloboryl ligands, BX_2 (X = F, Cl, Br, I), the earliest examples were reported by Nöth and Schmid in the 1960's.³ Thus in a series of papers, reviewed in ref. 3, these authors described a range of dichloroboryl species including [Co(BCl₂)(CO)₃(L)] (L = CO, PPh₃, AsPh₃), [Mn(BCl₂)(CO)₄(PPh₃)], [Fe(BCl₂)(CO)₂(η-C₅H₅)] and [Mo(BCl₂)(CO)₃(η-C₅H₅)] each derived from the reaction between BCl₃ and the corresponding metal carbonylate anion. Alternatively, dihaloboryl species were reportedly formed *via* B–X bond (X = halide) oxidative addition to Pt(0) or Co(0) centres, examples being the Pt(II) species [PtBr(BBr₂)(PPh₃)₂] and the Co(II), formally 19-electron, complexes [CoX(BX₂)(dppe)₂] (X = Cl, Br, I; dppe = 1,2-bis(diphenylphosphino)ethane),^{3,4} both proposed to have a *cis* configuration. The bis-boryl, 19-electron Co(II) complexes *trans*-[Co(BX₂)₂(dppe)₂] (X = Br, I) were also described,

University of Bristol, School of Chemistry, Bristol, UK BS8 1TS

derived from the reaction between [CoH(dppe)₂] and BX₃.^{3,5} These latter complexes were reported to be particularly versatile boryl transfer agents, the complex *trans*-[Co(BBr₂)₂(dppe)₂] being used to synthesise complexes of rhodium, copper and silver containing the BBr₂ ligand.^{3,5} Nevertheless, despite the impressive number of reports, none of these early compounds were structurally characterised and, partly on the basis of the ¹¹B NMR data presented, some doubt remains as to their correct formulation.

More recent work has provided the first definitive characterisation of dihaloboryl species. Thus, the first structurally characterised BF_2 complexes *cis*-[Pt(BF_2)₂(PPh₃)₂], $[Pt(BF_2)_2(dppb)](dppb = 1,2-bis(diphenylphosphino)butane)$ and fac-[Ir(BF₂)₃(CO)(PPh₃)₂] were recently described by us derived from the oxidative addition of B_2F_4 to Pt(0) and Ir(I) precursors respectively;6 spectroscopic data for an iridium BF₂ complex $[IrH(BF_2)(PMe_3)(\eta-C_5Me_5)]$ had previously been reported by Bergman and co-workers.⁷ In 2000, the osmium dichloroboryl complex $[OsCl(BCl_2)(CO)(PPh_3)_2]$ (δ_B 52.5) was described by Roper et al. having been prepared from the reaction between [OsCl(Ph)(CO)(PPh₃)₂] and HBCl₂.Et₂O.⁸ This and a number of subsequent reports revealed a rich chemistry of the BCl₂ ligand involving substitution at boron and resulting in a range of new boryl, tethered boryl and base-stabilised borylene complexes of osmium.^{8,9} In 2002, Aldridge reported spectroscopic data for the cyclopentadienyl complex [Fe(BCl₂)(CO)₂(η -C₅H₅)] ($\delta_{\rm B}$ 90.7),¹⁰ originally described by Nöth and Schmid,³ obtained from the reaction between BCl₃ and Na[Fe(CO)₂(η -C₅H₅)]. This was followed by a theoretical study on a range of similar metal boryl complexes, including BF₂ and BCl₂ examples, designed to probe the nature of the metal-boron bond.11 More recently, Braunschweig and co-workers have presented structural data for $[Fe(BCl_2)(CO)_2(\eta C_5H_5$] (δ_B 90.0), the first dichloroboryl complex to be structurally characterised.¹² These authors also reported the structure of the 4picoline adduct [Fe{BCl₂(4-pic)}(CO)₂(η -C₅H₅)](δ _B 18.8) in which the nitrogen atom is directly bonded to the boron centre.¹² In later reports, Braunschweig and co-workers have described the analogous pentamethylcyclopentadienyl complexes [Fe(BCl₂)(CO)₂(η- C_5Me_5] (δ_B 94.9)¹³ and [Fe(BBr₂)(CO)₂(η -C₅Me₅)] (δ_B 94),¹⁴ the former affording, on reaction with $[Pd(PCy_3)_2]$, the heteronuclear complex [$(\eta$ -C₅Me₅)Fe(μ -CO)₂(μ -BCl₂)Pd(PCy₃)] (δ _B 72.2) containing a bridging dichloroboryl ligand.13 Spectroscopic data were also presented for the phosphine adduct $[Fe{BCl_2(PCy_3)}(CO)_2(\eta C_5Me_5$] (δ_B 3.4).¹³ Both the cyclopentadienyl and pentamethylcyclopentadienyl iron dichloroboryl compounds have also been employed as precursors in the synthesis of dinuclear species containing substituent-free bridging boron atoms.15

In most of the recent studies, and much of the early work reported by Nöth and Schmid, the dihaloboryl species were prepared from a reaction between a boron trihalide and an anionic transition metal complex. The exceptions to this method involved an oxidative addition reaction either of a B–B bond, in the case of the BF₂ complexes,⁶ or of a boron–halogen bond in the case of some of the platinum and cobalt complexes described by Nöth and Schmid.³ In view of the known facile oxidative addition of B–X bonds to Pt(0), exemplified by the synthesis of such compounds as [PtCl{B(cat)}(PPh_3)₂],¹⁶ [PtCl{BCl(NMe₂)}(PPh_3)₂]¹⁷ and [PtBr{BBr(Fc)}(PCy_3)₂] (Fc = ferrocenyl; Cy = cyclohexyl)¹⁸ from ClB(cat), 1,2-B₂Cl₂(NMe₂)₂ and BBr₂(Fc) respectively, we were interested in establishing whether dihaloboryl compounds could be accessed directly from Pt(0) precursors and boron trihalides. Herein we describe the results of those studies.

Results and discussion

We were not the first to examine the reactions between phosphine-Pt(0) complexes and boron trihalides. Indeed, the reaction between BBr3 and [Pt(PPh3)4] in cyclohexane, reported by Nöth, Schmid and co-workers^{3,4} and mentioned above in the Introduction, was purported to give the Pt(II) complex $[PtBr(BBr_2)(PPh_3)_2]$. However, in subsequent work by Durkin and Schram¹⁹ and later Wallbridge and co-workers,20 compounds formulated as Lewis acid-base adducts were described. Thus Durkin and Schram reported that treatment of solid [Pt(PPh₃)₃] with gaseous BCl₃ afforded the 1: 2 adduct (Ph₃P)₃Pt·2BCl₃, the same species being formed, along with $Ph_3P \cdot BCl_3$, when $[Pt(PPh_3)_4]$ was employed.¹⁹ This 1:2 adduct was apparently unstable with respect to ligand dissociation when dissolved in a solvent and when the reaction between [Pt(PPh₃)₃] and BCl₃ was carried out in benzene solution, a product formulated as (Ph₃P)Pt·BCl₃ was proposed. In the later work by Wallbridge and co-workers,²⁰ the reaction between either $[Pt(PPh_3)_4]$ or $[Pt(PPh_3)_2(\eta-C_2H_4)]$ and BF₃ in toluene was reported to afford the adduct (Ph₃P)₂Pt·2BF₃ which was shown to react with BCl_3 to give $(Ph_3P)_2Pt \cdot 2BCl_3$.

In this study, the reaction between $[Pt(PMe_3)_4]$ (prepared *in situ* from $[Pt(nbe)_3]$ (nbe = norbornene) and four equivalents of PMe₃), and four equivalents of BCl₃ in toluene afforded the phosphine adduct Me₃PBCl₃ (δ_B 0.86, ¹ J_{PB} 164.7) and a platinum boryl species formulated as $[PtCl(BCl_2)(PMe_3)_2]$ (1), the ¹¹B NMR chemical shift and well resolved coupling to platinum being consistent

with the presence of a BCl₂ ligand ($\delta_{\rm B}$ 62.2, ¹ $J_{\rm PtB}$ 1214 Hz; $\delta_{\rm P}$ –14.6, ¹ $J_{\rm PtP}$ 2635 Hz). Attempts to crystallise 1 from this reaction mixture were unsuccessful, resulting only in small quantities of the species [PtCl(PMe₃)₃][BCl₄], although isolated yields of 1 as a white powder were typically >90%. [PtCl(PMe₃)₃][BCl₄] was characterised by X-ray crystallography, details for which are given in Table 1.

Similar reactions employing the phosphines PPh₃, PMePh₂, PMe_2Ph and PCy_3 (Cy = cyclohexyl) were also carried out. For PPh₃ and PMePh₂ the adducts Ph₃PBCl₃ and Ph₂MePBCl₃ were formed (the former characterised crystallographically) in addition to boryl products identified spectroscopically as $[PtCl(BCl_2)(PPh_3)_2]$ (2) $[\delta_B 59(br); \delta_P 23.0, {}^1J_{PtP} 3023 Hz]$ and $[PtCl(BCl_2)(PMePh_2)_2]$ (3) $[\delta_B \ 61(br); \ \delta_P \ 7.6, \ ^1J_{PtP} \ 2890 \ Hz]$ although in both these cases, the boryl signals in the ¹¹B NMR were broad and no coupling to platinum was resolved. Attempted crystallisation of 2 and 3 resulted only in the complexes $[PtCl(PPh_3)_3][BCl_4]$ and $[Pt_2(\mu-Cl)_2(PMePh_2)_4][BCl_4]_2$; the latter characterised by X-ray crystallography, details for which are given in Table 1. No spectroscopic evidence was observed for platinum boryl complexes when reactions involving the phosphines PMe₂Ph and PCy₃ were carried out, the only isolated products being PhMe₂PBCl₃ and $[Pt_2Cl_2(\mu-Cl)_2(PCy_3)_2]$ respectively, both identified by X-ray crystallography.

In all of the reactions carried out above, it was the tetraphosphine Pt(0) species, generated in situ, which was employed as the platinum starting material and BCl₃ was added in the molar ratio 4B : 1Pt. These reactions were attempted in light of some of the earlier studies referenced above but whilst they afforded the desired dichloroboryl species in some cases, the products were difficult to isolate not least because of the presence of the coformed phosphine-BCl₃ adducts. Reactions were therefore carried out in which only two equivalents of phosphine were added to the Pt(0) species $[Pt(nbe)_3]$ followed by one equivalent of BCl_3 . Thus reaction between [Pt(nbe)₃], two equivalents of the phosphines PPh₃, PMePh₂ or PMe₂Ph and one equivalent of BCl₃ afforded, in each case, high yields of the dichloroboryl species 2, 3 and $[PtCl(BCl_2)(PMe_2Ph)_2]$ (4) $[\delta_B = 61.8 (br); \delta_P - 6.7, {}^{1}J_{PtP} 2731 Hz].$ However, under similar conditions using PMe₃, although ³¹P NMR spectroscopy showed the presence of 1, the solutions decomposed rapidly to platinum metal. When an excess of BCl₃ was used, the unusual dinuclear species 5 was formed (see below).



2, R = Ph; **3**, $R_3 = Ph_2Me$; **4**, $R_3 = PhMe_2$

The complexes **2**, **3** and **4** were characterised by X-ray crystallography, the results for which are shown in Fig. 1–3; crystallographic data for all complexes are presented in Table 1. All three molecules (two per asymmetric unit for **4**) adopt similar structures in the solid state in which the boryl group is *trans* to the chloride, consistent with the known high *trans* influence of boryl ligands.^{1,21} The Pt–B bond distances (**2**, 1.972(3); **3**, 1.988(3); **4**, 1.963(6), 1.971(6) Å) are all similar but markedly shorter than the Pt–B distance in *trans*-[PtCl{B(cat)}(PPh₃)₂] (2.008(8) Å)¹⁶ although the

	[PtCl(PMe ₃) ₃][BCl ₄]] Ph ₃ PBCl ₃	$[Pt_2(\mu\text{-Cl})_2(PMePh_2)_4][BCl_4]_2$	PhMe ₂ PBCl ₃	$[Pt_2Cl_2(\mu-Cl)_2(PCy_3)_2]$	2	3
Formula M M Crystal system a/\hat{A} b/\hat{A} b/\hat{A} c/\hat{A} c/\hat{A} $a/^{\circ}$ $\beta/^{\circ}$ $\beta/^{\circ}$ $\gamma/^{\circ}$ Unit cell volume/ \hat{A}^{3} Space group Z μ/mm^{-1} Reflections measured/unique/observed ^a R_{1}^{h} (ϕ_{0}) wR_{2}° (ψ_{0}) wR_{2}° (ψ_{0})	$\begin{array}{c} C_9 H_{27} B Cl_5 P_3 Pt \\ 611.37 \\ Orthorhombic \\ 9.9079(4) \\ 13.2057(4) \\ 16.0316(6) \\ 16.0316(6) \\ 16.0316(6) \\ 2097.59(13) \\ P2_12_1 \\ 4 \\ 2097.59(13) \\ 875/3481/3297 \\ 6875/3481/3297 \\ 8.23 \\ 8.23 \\ 8.23 \end{array}$	C ₁₈ H ₁₅ BCl ₃ P 379.43 Monoclinic 11.8318(12) 9.7063(9) 15.8277(16) 92.467(2) 1816.0(3) P21/n 4 0.59 18842/4179/2819 6.26 4.19 12.25	$\begin{array}{c} C_{22}H_{32}B_{2}Cl_{10}P_{4}Pt_{2}\\ 1567.12\\ 7567.12\\ 1567.12\\ 1567.12\\ 9.896(2)\\ 11.707(2)\\ 11.707(2)\\ 12.654(3)\\ 84.45(3)\\ 84.45(3)\\ 79.32(3)\\ 84.45(3)\\ 12.654(3)\\ 79.32(3)\\ 79.32(3)\\ 79.32(3)\\ 79.32(3)\\ 14.31.6(5)\\ 11.70(5)\\ 12.557$	C ₈ H ₁ BCl ₃ P 255.30 Orthorhombic 6.2473(7) 10.8165(11) 17.5194(18) 17.5194($C_{43}H_{44}CI_{4}P_{2}Pt_{2}$ 1184.94 Monoclinic 14.229(3) 16.041(3) 19.920(4) 97.22(3) 4510.5(16) C2/c 6.53 25604/5181/4901 3.13 1.98 4.74	C ₃₆ H ₃₀ BCl ₃ P ₂ Pt 836.79 Monoclinic 28.471(6) 11.886(2) 19.635(4) 100.45(3) 6534(2) 6534($\begin{array}{c} C_{26}H_{26}BCl_3P_2Pt\\ 712.66\\ Monoclinic\\ 14.938(3)\\ 10.071(2)\\ 18.279(4)\\ 101.18(3)\\ 2697.6(10)\\ P2_1/n\\ P2_1/n\\ 2563\\ 30316/6189/5803\\ 30316/6189/5803\\ 6.35\\ 6.35\end{array}$
	4	$Cl_3 BP(Ph)_2 CH_2(Ph)_2 PBCl_3$	5	$[Pt(PMe_3)_2(dppm)]Cl_2$	8	10	11
Formula M M Crystal system a/\hat{A} b/\hat{A} b/\hat{A} b/\hat{A} c/\hat{A} $\beta/^{\circ}$ Unit cell volume/ \hat{A}^3 $g/^{\circ}$ Unit cell volume/ \hat{A}^3 $g/^{\circ}$ $g/^{\circ}$ $M^{\rm IIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIII$	$\begin{array}{c} C_{16}H_{22}BCl_3P_2Pt\\ 588.53\\ Monoclinic\\ 17.466(4)\\ 6.0751(12)\\ 0.0751(12)\\ 21.219(4)\\ 113.81(3)\\ 2059.9(9)\\ 113.81(3)\\ 2059.9(9)\\ 113.81(3)\\ 2059.9(9)\\ 20$	$\begin{array}{c} C_{25}H_{22}B_{2}Cl_{6}P_{2}\\ 618.68\\ Monoclinic\\ 11.066(2)\\ 10.003(2)\\ 10.003(2)\\ 10.003(2)\\ 129.95(2)\\ 129.95(2)\\ 1389.2(7)\\ Pc\\ Pc\\ Pc\\ 1389.2(7)\\ 1389.2(7)\\ 1389.2(7)\\ 0.03(4)\\ 0.05(4)\\ 0.05(4)\\ 0.05(4)\end{array}$	$C_{12}H_{36}B_{3}Cl_{9}P_{4}Pt_{2}$ 1045.95 Monoclinic 12.306(3) 13.214(3) 20.313(4) 93.35(3) 3297.5(13) $P2_{1}/n$ 9.40 36989/7566/6902 3.60 5.00 2.48 5.00	C ₃₁ H ₄₀ Cl ₂ P ₄ Pt 802.50 Orthorhombic 29.9175(16) 20.6730(11) 11.5155(6) 7122.2(7) <i>Pbcn</i> 8 8 4.29 71496/8138/6996 7.24 7.24	C ₁₀ H ₃₈ BCl ₂ NP ₂ Pt 501.07 Orthorhombic 10.125(2) 11.154(2) 1852.6(6) P2 ₁ 2 ₁ 2 ₁ 8.02 21217/4242/4234 2.85 1.90 4.93 0.004(6)	$\begin{array}{c} C_{12}H_{22}BCIO_2P_2Pt\\ 501.59\\ Monoclinic\\ 5.8930(16)\\ 22.748(9)\\ 12.946(4)\\ 95.56(3)\\ 1122.46(4)\\ 95.56(3)\\ 1127.3(10)\\ P2_{1}/c\\ 4\\ 1\\ 18.33\\ 12164/2972/2858\\ 6.13\\ 6.13\\ 6.13\\ 6.13\\ 6.13\end{array}$	$C_{18}H_{26}B_2O_4P_2Pt$ 585.04 Monoclinic 14.4914(3) 9.3235(2) 16.1496(3) 90.4770(10) 2181.91(8) $P2_1/n$ 13.58 16409/3893/3792 5.95 5.95 7.67

Table 1 Crystallographic data





Fig. 1 A view of the molecular structure of compound 2. Selected bond length (Å) and angle (°) data include: Pt(1)-B(1) 1.972(3), Pt(1)-Cl(1) 2.4368(8), Pt(1)-P(1) 2.3005(7), Pt(1)-P(2) 2.2950(7), B(1)-Cl(2) 1.791(3), B(1)-Cl(3) 1.779(3); Cl(1)-Pt(1)-B(1) 173.03(8), P(1)-Pt(1)-P(2) 169.45(2), Cl(2)-B(1)-Cl(3) 113.12(4), Pt(1)-B(1)-Cl(2) 119.85(14), Pt(1)-B(1)-Cl(3) 127.03(15); interplanar angle 81.0.



Fig. 2 A view of the molecular structure of compound 3. Selected bond length (Å) and angle (°) data include: Pt(1)-B(1) 1.988(3), Pt(1)-Cl(1) 2.4461(8), Pt(1)-P(1) 2.2961(9), Pt(1)-P(2) 2.3005(9), B(1)-Cl(2) 1.789(3), B(1)-Cl(3) 1.757(3); Cl(1)-Pt(1)-B(1) 173.25(9), P(1)-Pt(1)-P(2) 173.51(2), Cl(2)-B(1)-Cl(3) 113.57(18), Pt(1)-B(1)-Cl(2) 119.16(18), Pt(1)-B(1)-Cl(3) 127.27(17); interplanar angle 92.7.

This journal is \odot The Royal Society of Chemistry 2007

Fig. 3 A view of the molecular structure of compound **4**. Selected bond length (Å) and angle (°) data include: Molecule 1 Pt(1)–B(1) 1.963(6), Pt(1)–Cl(1) 2.4385(15), Pt(1)–P(1) 2.3065(15), Pt(1)–P(2) 2.3003(15), B(1)–Cl(2) 1.794(7), B(1)–Cl(3) 1.778(7); Cl(1)–Pt(1)–B(1) 176.8(2), P(1)–Pt(1)–P(2) 176.22(5), Cl(2)–B(1)–Cl(3) 111.2(3), Pt(1)–B(1)–Cl(2) 124.9(4), Pt(1)–B(1)–Cl(3) 123.9(4); interplanar angle 85.8. Molecule 2 Pt(2)–B(2) 1.971(6), Pt(2)–Cl(4) 2.4433(15), Pt(2)–P(3) 2.3017(15), Pt(2)–P(4) 2.2994(15), B(2)–Cl(5) 1.780(7), B(2)–Cl(6) 1.782(7); Cl(4)–Pt(2)–B(2) 173.3(2), P(3)–Pt(2)–P(4) 175.46(5), Cl(5)–B(2)–Cl(6) 111.4(3), Pt(2)–B(2)–Cl(5) 125.1(4), Pt(2)–B(2)–Cl(6) 123.5(4); interplanar angle 92.8.

Pt-Cl distances are more comparable (2, 2.4368(8); 3, 2.4461(8); 4, 2.4385(15), 2.4433(15); $[PtCl{B(cat)}(PPh_3)_2]$ 2.4446(14) Å).¹⁶ In the complex trans-[PtCl{BCl(NMe₂)}(PPh₃)₂],¹⁷ the Pt-B and Pt-Cl distances are 2.075(10) and 2.470(2) Å, respectively, revealing that the Pt-B bond length is broadly correlated with the expected Lewis acidity of the boryl group, i.e. shorter for a more Lewis acidic boron. Each boron centre in 2, 3 and 4 is trigonal planar, the Cl-B-Cl angle being a little less than 120° and the two Pt-B-Cl angles being slightly greater (the slight asymmetry in 2 and 3 is not present in 4); a similar set of angles is observed for $[Fe(BCl_2)(CO)_2(\eta-C_5H_5)]$.¹² In all cases, the angle between the mean platinum square plane and the mean boron trigonal plane is close to 90° (2, 81.0; 3, 92.7; 4, 85.8, 92.8°) which compares with the corresponding angles for $[PtCl{B(cat)}(PPh_3)_2]$ (78.3°)¹⁶ and [PtCl{BCl(NMe₂)}(PPh₃)₂] (90.0°).¹⁷ We note a recent report from Marder and co-workers²¹ describing DFT studies on Pt(II) boryl species of the type $[PtCl(BR_2)(PMe_3)_2]$. The trends observed experimentally in the Pt-B and Pt-Cl bond lengths are in full accord with those predicted by the DFT calculations. Thus the trans influence of the B(cat) and BCl₂ groups was calculated to be nearly identical, whilst shorter Pt-B distances were anticipated for Pt-BCl₂ (1.976 Å) vs. Pt-B(cat) (2.005 Å).²¹

Reactions involving the diphosphines dppe (1,2-bis(diphenylphosphino)ethane) and dppm (bis(diphenylphosphino)methane) were also carried out. Thus reaction between $[Pt_2(dppm)_3]$ and BCl₃ afforded a mixture of products, one of which was shown by X-ray crystallography to be the dppm–BCl₃ adduct $Cl_3BP(Ph)_2CH_2(Ph)_2PBCl_3$, details of which are given in Table 1.

Crystals of the unusual dinuclear species $[Pt_2(BCl_2)_2(PMe_3)_4(\mu-Cl)][BCl_4]$ (5) were obtained. A view of the structure of the cation is shown in Fig. 4 which reveals an arrangement in which two square planar platinum(II) centres are linked by a single bridging chloride which is *trans* to a BCl₂ group on each platinum centre. Both Pt–B bonds [1.960(5) and 1.955(4) Å] are similar in length to those observed for **2**, **3** and **4** as are the Pt–Cl bond lengths [2.5155(10) and 2.5088(11) Å]; the boron trigonal planes are also close to orthogonal with respect to the platinum square planes (89.4 and



Fig. 4 A view of the molecular structure of compound **5**. Selected bond length (Å) and angle (°) data include: Pt(1)-B(1) 1.960(5), Pt(2)-B(2) 1.955(4), Pt(1)-Cl(3) 2.5155(10), Pt(2)-Cl(3) 2.5088(11), B(1)-Cl(1) 1.759(5), B(1)-Cl(2) 1.778(5), B(2)-Cl(4) 1.780(5), B(2)-Cl(5) 1.771(5); B(1)-Pt(1)-Cl(3) 175.54(13), P(1)-Pt(1)-P(2) 176.11(4), B(2)-Pt(2)-Cl(3) 171.40(13), P(3)-Pt(2)-P(4) 168.70(4), Pt(1)-Cl(3) 126.48(4); interplanar angles 89.4 [B(1)] and 86.3 [B(2)].

 86.3°). Whilst the mechanism by which compound **5** is formed is unclear, a formal equation for its formation, eqn (1), is consistent with the expected high *trans* influence of the boryl groups resulting in the lability of a *trans* chloride which is subsequently scavenged by BCl₃.



Having a reliable route to quantities of $[PtCl(BCl_2)(PMe_3)_2]$ (1), enabled a study of its reactivity to be carried out which is summarised in Scheme 1. Thus treatment of 1 with NEt₃ or pyridine (py) afforded the adducts $[PtCl{BCl_2(NEt_3)}(PMe_3)_2]$ (6) $(\delta_B 27.8; \delta_P - 13.0, {}^1J_{PtP} 2643 \text{ Hz})$ and $[PtCl{BCl_2(py)}(PMe_3)_2]$ (7) $(\delta_B 29.9, {}^1J_{PtB} 957; \delta_P - 10.4, {}^1J_{PtP} 2694 \text{ Hz})$ respectively, both characterised spectroscopically. Proposed structures for 6 and 7 are shown in the diagrams below consistent both with the ${}^{11}B$ NMR data and the known structure of the species $[Fe{BCl_2(4-pic)}(CO)_2(\eta-C_5H_5)].{}^{12}$ Treatment of 1 with dppm afforded a mixture of products of which only the salt $[Pt(PMe_3)_2(dppm)]Cl_2$ was characterised; X-ray crystallographic data for this species are given in the Experimental section.





The reaction between 1 and either four equivalents of NHEt₂ or piperidine (pipH) resulted in the mono-substituted boryl species *trans*-[PtCl{BCl(NEt₂)}(PMe₃)₂] (8) ($\delta_{\rm B}$ 35.9, ¹ $J_{\rm PtB}$ 1028; $\delta_{\rm P}$ 13.1, ${}^{1}J_{\text{PtP}}$ 2772 Hz) and [PtCl{BCl(pip)}(PMe_{3})_{2}] (9) (δ_{B} 34.3; δ_{P} -9.9, ${}^{1}J_{PtP}$ 3000 Hz) respectively, the eliminated HCl in each case removed as [NH₂Et₂]Cl or [pipH₂]Cl. Even with sufficient diethylamine or piperidine present, no evidence was obtained for a disubstituted boryl species. Spectroscopic data for 8 and 9 were consistent with the proposed structures and compound 8 was also characterised by X-ray crystallography, the results of which are shown in Fig. 5. The structure of 8 is similar to those of 2, 3 and 4 described above. Notable differences, however, are the longer Pt-B and Pt-Cl bond lengths [2.005(4) and 2.4604(10) Å respectively], the former a likely consequence of the less Lewis acidic boron centre resulting from the π -donor diethylamido substituent as described earlier. The angle between the platinum mean square plane and the boron mean trigonal plane is 90.0° and that between the adjacent boron and nitrogen trigonal planes is 6.0° (consistent with the π -donor properties of the amido nitrogen). In all respects, the structure of 8 is similar to that previously reported for the dimethylamido analogue trans-[PtCl{BCl(NMe₂)}(PPh₃)₂].¹⁷



8, X = NEt₂; **9**, X = pip

Treatment of either **1** (in the presence of excess NEt₃) or **8** with catechol afforded the B(cat) derivative *trans*-[PtCl{B(cat)}(PMe₃)₂] (**10**) (Schemes 1 and 2) characterised spectroscopically (δ_B 33.3, ¹ J_{PtB} 1039; δ_P –12.6, ¹ J_{PtP} 2768 Hz) and by



Fig. 5 A view of the molecular structure of compound 8. Selected bond length (Å) and angle (°) data include: Pt(1)-B(1) 2.005(4), Pt(1)-Cl(1) 2.4604(10), Pt(1)-P(1) 2.2835(10), Pt(1)-P(2) 2.2795(10), B(1)-Cl(2) 1.843(5), B(1)-N(1) 1.394(6); Cl(1)-Pt(1)-B(1) 178.86(15), P(1)-Pt(1)-P(2) 178.33(4), Cl(2)-B(1)-N(1) 115.5(3), Pt(1)-B(1)-Cl(2) 117.1(2), Pt(1)-B(1)-N(1) 127.4(3); interplanar angle 90.0.

X-ray crystallography. The results of the structure determination are shown in Fig. 6 and reveal the expected *trans* geometry as observed for related species including the chloro–B(cat) complex *trans*-[PtCl{B(cat)}(PPh_3)₂].¹⁶ As shown in Scheme 2, complex **10** was formed in the reaction between [Pt(PMe₃)₄] and ClB(cat) and also from the slow decomposition of *cis*-[Pt{B(cat)}₂(PMe₃)₂] (**11**) in dichloromethane over a period of months. Compound **11** ($\delta_{\rm B}$ 49.4; $\delta_{\rm P}$ –14.3, ¹ $J_{\rm PtP}$ 1490 Hz) was prepared from the reaction between [Pt(PMe₃)₄] and B₂(cat)₂, by analogy with other Pt(II) bisboryl species,^{1,16,17,22} and is structurally similar (Fig. 7) to all related bis-phosphine, platinum(II) bis-boryl species.^{1,16,17,22}

Me₃P

PMe₃

B(cat)

11

-B(cat)

PMe₃

. PMe₃ -B(cat)





Fig. 6 A view of the molecular structure of compound 10. Selected bond length (Å) and angle (°) data include: Pt(1)-B(1) 1.995(6), Pt(1)-Cl(1) 2.4308(16), Pt(1)-P(1) 2.284(2), Pt(1)-P(2) 2.289(2), B(1)-O(1) 1.423(7), B(1)-O(2) 1.415(7); Cl(1)-Pt(1)-B(1) 178.15(18), P(1)-Pt(1)-P(2) 178.48(5), O(1)-B(1)-O(2) 108.8(5), Pt(1)-B(1)-O(1) 124.2(4), Pt(1)-B(1)-O(2) 127.0(4); interplanar angle 95.8.



Fig. 7 A view of the molecular structure of compound 11. Selected bond length (Å) and angle (°) data include: Pt(1)-B(1) 2.061(10), Pt(1)-B(2) 2.032(9), Pt(1)-P(1) 2.331(2), Pt(1)-P(2) 2.327(2), B(1)-O(1) 1.413(12), B(1)-O(2) 1.388(14), B(2)-O(3) 1.415(11), B(2)-O(4) 1.413(11); B(1)-Pt(1)-B(2) 82.2(2), P(1)-Pt(1)-P(2) 101.47(8), O(1)-B(1)-O(2) 108.1(7), Pt(1)-B(1)-O(1) 126.7(7), Pt(1)-B(1)-O(2) 124.6(7), O(3)-B(2)-O(4) 107.8(7), Pt(1)-B(2)-O(3) 128.6(7), Pt(1)-B(2)-O(4) 123.5(6); interplanar angles 81.4 [B(1)], 91.5 [B(2)].

In conclusion, these studies have shown that platinum(II) dichloroboryl complexes of the form *trans*-[PtCl(BCl₂)(PR₃)₂] can be formed in high yield from the oxidative addition of a B–Cl bond in BCl₃ to a platinum(0) centre in the presence of two equivalents of a phosphine ligand. Furthermore, an examination of the reactivity of such species, *trans*-[PtCl(BCl₂)(PMe₃)₂] in particular, reveals an adduct chemistry associated with the Lewis acidic boron

centre, consistent with the previous work of Braunschweig,¹² and a substitution chemistry of the B–Cl bond(s) which parallels the chemistry observed by Roper for the related osmium BCl₂ complex described in the Introduction.^{8,9} The original report by Nöth and Schmid of oxidative addition of BBr₃ to Pt(0) resulting in a BBr₂ ligand is therefore consistent with these studies. In contrast, however, we obtained no evidence for BCl₃ adducts of platinum(0) as described by Schram¹⁹ and later Wallbridge.²⁰

Experimental

General procedures

All reactions were performed under dry dinitrogen using standard Schlenk line techniques. Dry solvents were obtained from an anhydrous engineering system and were degassed prior to use. All reagents were procured commercially and used without further purification except for $[Pt(nbe)_3]^{23}$ (nbe = norbornene = bicy-clo[2.2.1]heptene) and B₂(cat)₂²⁴ which were prepared by literature methods.

Preparations

[PtCl(BCl₂)(PMe₃)₂] (1). [Pt(nbe)₃] (0.10 g. 0.21 mmol) was dissolved in hexane (5 cm^3) , to which PMe₃ $(0.099 \text{ cm}^3, 0.84 \text{ mmol})$ was added through a micro-syringe. The reaction mixture was stirred for 1 h, and the solvent was removed in vacuo to give a white solid. The solid was redissolved in toluene (10 cm³), and BCl₃ (0.84 cm³, 0.84 mmol, 1 M in heptane) was added at -78 °C. The mixture was allowed to warm to room temperature, and stirred for 1 h during which time a quantity of white precipitate formed. After this time, all volatiles were removed in vacuo, and the solid was redissolved in toluene (10 cm³) and filtered, affording a colourless solution which was allowed to stand overnight resulting in a further precipitate of the by-product, Me₃PBCl₃. The solvent was removed slowly in vacuo, and the resulting solid redissolved in toluene (5 cm^3), and filtered to produce a solution of 1 from which a white powder was obtained after further solvent removal (0.096 g, 97%). Spectroscopic data for 1: NMR $(C_6D_6)^{11}B{^1H}$ δ 62.2 (¹J_{PtB} 1214 Hz), ³¹P{¹H} δ –14.6 (¹J_{PtP} 2635 Hz); ¹H δ 1.12 (m), ${}^{13}C{}^{1}H{}\delta$ 13.9 (t, ${}^{1}J_{PC}$ 20 Hz); mass spectrum (EI) 383 $(M^+ - BCl_2)$. Attempts to crystallise 1 from this reaction mixture were unsuccessful, resulting only in small quantities of the species identified as [PtCl(PMe₃)₃][BCl₄] by X-ray crystallography.

[PtCl(BCl₂)(PPh₃)₂] (2). [Pt(nbe)₃] (0.20 g. 0.42 mmol) and PPh₃ (0.22 g, 0.84 mmol) were dissolved in toluene (5 cm³) and the reaction mixture stirred for 1 h, after which time the solvent was removed *in vacuo* to give a white solid. This solid was then dissolved in toluene (5 cm³), and BCl₃ (0.42 cm³, 0.42 mmol, 1 M in heptane) was added at -78 °C. The mixture was allowed to warm to room temperature, and stirred for 1 h during which time a white precipitate formed. All volatiles were then removed *in vacuo*, and the resulting solid washed with toluene (3 × 1 cm³) (0.29 g, 81%). Colourless crystals suitable for X-ray crystallography were formed from solutions in toluene (1 cm³) and CH₂Cl₂ (1 cm³) layered with hexane. Spectroscopic data for **2**: NMR (C₆D₆) ¹¹B{¹H} δ 59 (br), ³¹P{¹H} δ 23.0 (¹J_{PLP} 3023 Hz), ¹H δ 7.66–7.73 (m), 7.33–7.41 (m), ¹³C{¹H} δ 140.0 (s), 134.9 (t, J_{PC} 6.0), 130.6 (s), 128.3 (t, J_{PC} 4.5); mass spectrum (EI) 754 (M⁺ – BCl₂). Anal. Calcd for $C_{36}H_{30}BCl_3P_2Pt$: H, 3.60; C, 51.65%. Found H, 3.70; C, 51.60%.

[PtCl(BCl₂)(PMePh₂)₂] (3). [Pt(nbe)₃] (0.10 g. 0.21 mmol) and PMePh₂ (0.080 cm³, 0.42 mmol) were dissolved in toluene (5 cm³) and the reaction mixture was stirred for 1 h, after which time the solvent was removed *in vacuo* to give an off-white oil. This oil was then dissolved in toluene (5 cm³), and BCl₃ (0.21 cm³, 0.21 mmol, 1 M in heptane) was added at -78 °C. The mixture was allowed to warm to room temperature, and stirred for 1 h. All volatiles were then removed *in vacuo*, and the resulting solid was redissolved in toluene (3 cm³) and filtered. The filtrate was evaporated *in vacuo* to give a white solid (0.085 g, 57%). Colourless crystals suitable for X-ray crystallography were formed from solutions in toluene (1 cm³) layered with hexane. Spectroscopic data for **3**: NMR (C₆D₆) ¹¹B{¹H} δ 61 (br), ³¹P{¹H} δ 7.6 (¹J_{PIP} 2890 Hz); mass spectrum (EI) 631 (M⁺ – BCl₂).

 $[PtCl(BCl_2)(PMe_2Ph)_2]$ (4). $[Pt(nbe)_3]$ (0.10 g. 0.21 mmol) and PMe₂Ph (0.061 cm³, 0.42 mmol) were dissolved in toluene (5 cm³) and the reaction mixture stirred for 1 h, after which time the solvent was removed in vacuo to give an off-white oil. This oil was then dissolved in toluene (5 cm³), and BCl₃ (0.21 cm³, 0.21 mmol, 1 M in heptane) was added at -78 °C. The mixture was allowed to warm to room temperature, and stirred for 1 h. All volatiles were then removed in vacuo, and the resulting solid was redissolved in toluene (3 cm³) and filtered. The filtrate was evaporated in vacuo to give a white solid (0.087 g, 74%). Colourless crystals suitable for X-ray crystallography were formed from solutions in toluene (1 cm³) layered with hexane. Spectroscopic data for 4: NMR $(C_6 D_6)^{11}B{}^{1}H{}\delta 61.8$ (br), ${}^{31}P{}^{1}H{}\delta -6.7$ (${}^{1}J_{PtP}$ 2731 Hz), ${}^{1}\text{H}\,\delta$ 7.58–7.65 (m), 7.00–7.08 (m), 1.56 (t, ${}^{3}J_{\text{PtH}}$ 18.0, ${}^{2}J_{\text{PH}}$ 3.0 Hz), $^{13}C{^{1}H} \delta$ 131.3 (t, J_{PC} 6.0), 130.1 (s), 128.4 (t, J_{PC} 5.3), 13.3 (t, J_{PC} 19.6); mass spectrum (EI) 507 (M⁺ – BCl₂). Anal. Calcd for C16H22BCl3P2Pt: H, 3.75; C, 32.65%. Found H, 3.85; C, 32.75%.

 $[Pt_2(BCl_2)_2(PMe_3)_4(\mu-Cl)][BCl_4]$ (5). $[Pt(nbe)_3]$ (0.10 g. 0.21 mmol) was dissolved in hexane (5 cm³), to which PMe₃ (0.055 cm³, 0.42 mmol) was added through a micro-syringe. The reaction mixture was stirred for 1 h, and all the volatiles were removed in vacuo to give an off-white oil. The solid was redissolved in toluene (5 cm³), and an excess of BCl₃ (0.4 cm³, 0.46 mmol, 1 M in heptane) was added at -78 °C. The mixture was allowed to warm to room temperature, and stirred for 1 h during which time a quantity of black oil formed. After this time, all volatiles were removed in vacuo, and the solid was redissolved in toluene (10 cm³) and filtered, affording a colourless solution. The solvent was removed slowly in vacuo to give a white powder (0.047 g, 43%). Colourless crystals suitable for X-ray crystallography were formed from solutions in toluene (1 cm³) layered with hexane. Spectroscopic data for 5: NMR (C_6D_6) ¹¹B{¹H} δ 60.6 (¹J_{PIB} 1242 Hz), 11.1 (s), ³¹P{¹H} δ -14.7 (¹J_{PIP} 2628 Hz), ¹H δ 1.08 (t, ³J_{PtH} 18.0, ²J_{PH} 3.0 Hz), ¹³C{¹H} δ 14.0 (t, J_{PC} 19.6 Hz); mass spectrum (EI) 383 (M⁺ – Pt(BCl₂)₂). Anal. Calcd for C₁₂H₃₆B₃Cl₉P₄Pt₂: H, 3.45; C, 13.80%. Found H, 3.40; C, 13.80%.

 $[PtCl{BCl_2(NEt_3)}(PMe_3)_2]$ (6). A sample of compound 1 (0.10 g, 0.21 mmol) was dissolved in toluene (5 cm³), to which an

excess of NEt₃ (0.5 cm³) was then added. The mixture was stirred overnight affording a white precipitate of compound **6** (0.043 g, 36.9%). Spectroscopic data for **6**: NMR (CD₃CN) ¹¹B{¹H} δ 27.8 (br), ³¹P{¹H} δ -13.0 (¹*J*_{PIP} 2643 Hz), ¹H δ 3.58–3.65 (q), 1.88 (t), 1.21 (br).

[PtCl{BCl₂(py)}(PMe₃)₂] (7). A sample of compound 1 (0.10 g, 0.21 mmol) was dissolved in toluene (5 cm³) and the solution was layered with an excess of pyridine (3 cm³). After one week, a white precipitate of compound 7 was obtained which was washed with hexane (3 × 2 cm³) (0.052 g, 46.5%). Spectroscopic data for 7: NMR (CD₃CN) ¹¹B{¹H} δ 29.9 (¹*J*_{PtB} 957 Hz), ³¹P{¹H} δ -10.4 (¹*J*_{PtP} 2694 Hz). Both 6 and 7 are prone to loss of the Lewis base bound to boron which prevented more extensive spectroscopic and analytical characterisation.

 $[PtCl\{BCl(NEt_2)\}(PMe_3)_2]$ (8). A sample of compound 1 (0.10 g, 0.21 mmol) was dissolved in toluene (10 cm^3) , to which HNEt₂ (0.085 cm³, 0.84 mmol) was added. After stirring for 2 h, the solution become cloudy and a white precipitate formed. The solution was filtered and allowed to stand overnight affording a precipitate of [NH₂Me₂]Cl. After filtration and removal of the solvent, compound 8 was obtained as a white solid (0.095 g,91.7%). Colourless crystals suitable for X-ray crystallography were formed from solutions in toluene (0.5 cm³) layered with hexane. Spectroscopic data for 8: NMR (C_6D_6) ¹¹B{¹H} δ 35.9 (¹ J_{PtB} 1028 Hz), ${}^{31}P{}^{1}H{}\delta -13.1 ({}^{1}J_{PtP} 2772 \text{ Hz}), {}^{1}H \delta 3.41-3.35 (q, 2H);$ 3.24–3.19 (q, 2H,); 1.26 (t, 6H, ³*J*_{PtH} 12.0, ²*J*_{PH} 3.0 Hz); 0.95 (t, 3H); 0.88 (t, 3H), ${}^{13}C{}^{1}H{}\delta$ 47.5 (CH₂, ${}^{3}J_{PtC}$ 70.4 Hz), 41.3 (CH₂, ${}^{3}J_{PtC}$ 22.1 Hz), 14.7 (s, CH₃), 14.5 (m, ${}^{1}J_{PC}$ 19.1, ${}^{2}J_{PtC}$ 38.2 Hz); mass spectrum (EI) 465(M^+ - Cl). Anal. Calcd for $C_{10}H_{28}BCl_2NP_2Pt{\rm :}$ H, 5.65; C, 23.95%. Found H, 5.60; C, 24.00%.

[PtCl{BCl(pip)}(PMe_3)2] (9). A sample of compound 1 (0.10 g, 0.21 mmol) was dissolved in toluene (10 cm³), to which piperidine (0.081 cm³, 0.84 mmol) was added. After stirring for 2 h, the solution become cloudy and a white precipitate formed. The solution was filtered and allowed to stand affording a precipitate of [pipH₂]Cl. After filtration and removal of the solvent, compound 9 was obtained as a white solid (0.083 g, 78.3%). Spectroscopic data for 9: NMR (C_6D_6) ¹¹B{¹H} δ 34.3 (br), ³¹P{¹H} δ –9.9 (¹*J*_{PtP} 3000 Hz). Despite the similarity to compound **8**, which has been fully characterised, satisfactory analytical data were not obtained for **9**.

[PtCl{B(cat)}}(PMe₃)₂] (10). A sample of compound 1 (0.10 g, 0.21 mmol) was dissolved in toluene (5 cm³), and an excess of NEt₃ (0.5 cm³) was added followed by a solution of catechol (0.023 g, 0.21 mmol) in toluene (5 cm³). The mixture was stirred for 2 h at room temperature affording compound **10** as a white solid (0.076 g, 72%). Spectroscopic data for **10**: NMR (CH₃CN)¹¹B{¹H} δ 33.3 (¹J_{PtB} 1039 Hz), ³¹P{¹H} δ -12.6 (¹J_{PtP} 2768 Hz), ¹⁴ δ 7.15–7.25 (m, C₆H₄); 121.7 (s, C₆H₄); 111.4 (m, CH₃), ¹³C{¹H} δ 149.9 (s, C₆H₄); 121.7 (s, C₆H₄); 111.4 (s, C₆H₄); 14.8 (t, CH₃, ¹J_{PC} 20.1); mass spectrum (EI) 383 (M⁺ – Bcat). Anal. Calcd for C₁₂H₂₂BClO₂P₂Pt: H, 4.40; C, 28.75%. Found H, 4.30; C, 28.70%. Compound **10** was also prepared, although not isolated, by the following methods and identified spectroscopically (¹¹B and ³¹P NMR).

Method (i). A sample of compound **8** (0.10 g, 0.19 mmol) was dissolved in toluene (5 cm³) to which a solution of catechol (0.063 g, 0.57 mmol) in toluene (5 cm³) was added.

Method (*ii*). A sample of compound **11** (0.11 g, 0.18 mmol) was dissolved in CH_2Cl_2 (5 cm³) and the solution allowed to stand for 2 months. After this time, colourless crystals of **10** were obtained by layered the solution with hexane.

Method (*iii*). A sample of $[Pt(PMe_3)_4]$ (0.11 g, 0.21 mmol) was dissolved in toluene (5 cm³) to which a solution of ClB(cat) (32 mg, 0.21 mmol) in toluene (5 cm³) was added.

 $[Pt{B(cat)}_2(PMe_3)_2]$ (11). A sample of $[Pt(nbe)_3]$ (0.10 g, 0.21 mmol) was dissolved in hexane (5 cm³), to which PMe₃ (0.099 cm³, 0.84 mmol) was added through a micro-syringe. The reaction mixture was stirred for 1 h at room temperature and the solvent removed in vacuo affording a white solid. This solid was dissolved in toluene (10 cm³), and B₂(cat)₂ (0.05 g, 0.21 mmol) was added. The resulting mixture was stirred for 2 h at room temperature to give white precipitate. The solvent volume to was reduced to 1.5 cm³ in vacuo and the white precipitate was filtered off, washed with toluene (2 cm³) and hexane (2 \times 2 cm³) and dried in vacuo (0.081 g, 65%). Colourless crystals suitable for Xray crystallography were formed from solutions in CH_2Cl_2 (1 cm³) layered with hexane. Spectroscopic data for 11: NMR (CD₂Cl₂) ¹¹B{¹H} δ 49.4 (br), ³¹P{¹H} δ -14.3 (¹*J*_{PtP} 1490 Hz), ¹H δ 7.13– 7.07 (m, C₆H₄, 4H), 6.91–6.86 (m, C₆H₄, 4H), 1.50 (br, CH₃, 18H), $^{13}C{^{1}H}\delta$ 149.9 (s, C₆H₄), 120.8 (s, C₆H₄), 111.4 (s, C₆H₄), 19.2 (t, CH₃, ${}^{1}J_{PC}$ 12.1); mass spectrum (EI) 346 (M⁺ – 2Bcat). Anal. Calcd for C₁₈H₂₆B₂O₄P₂Pt: H, 4.50; C, 36.95%, Found H, 4.40; C, 37.00%.

X-Ray crystallography

Crystallographic data for compounds Ph₃BCl₃, PhMe₂PBCl₃, 3, 4 and Cl₃BP(Ph)₂CH₂(Ph)₂PBCl₃, were collected at 173 K on a Bruker-AXS SMART 1K diffractometer using Mo-Ka radiation $(\lambda = 0.71073 \text{ Å})$. For compounds $[Pt_2(\mu-Cl)_2(PMePh_2)_4][BCl_4]_2$, $[Pt_2Cl_2(\mu-Cl)_2(PCy_3)_2]$, 2 and 8, data were collected at 100 K on a Bruker-AXS SMART Apex diffractometer equipped with monocapillary beam focussing optics using Mo-Ka radiation $(\lambda = 0.71073 \text{ Å})$. For compounds [PtCl(PMe₃)₃][BCl₄], 10 and 11, data were collected using Cu-Ka radiation ($\lambda = 1.5418$ Å) at 100 K on a Bruker-AXS Proteum diffractometer equipped with a rotating anode source operated at 4.8 kW and Osmic CMF12-38Cu6 (blue) mirror optics. For all compounds, intensity data were collected as a series of frames, each of ω width 0.3°, integrated²⁵ and corrected for absorption²⁶ and solved and refined using routine techniques.²⁷ The symmetrically unique anion in complex $[Pt_2(\mu-Cl)_2(PMePh_2)_4][BCl_4]_2$ shows positional disorder for three of its chlorine atoms. A sensible model for this disorder was obtained by restraining the sets of $Cl \cdots Cl$ distances in each of the disordered components to be equal and restraining the displacement parameters of pairs of chlorine atoms opposite each other with respect to the undisordered B(1)–Cl(2) axis to be equal. Attempts to find a satisfactory model for the solvent present in the structure of [Pt(PMe₃)₂(dppm)]Cl₂ were unsuccessful and a diffuse solvent correction was applied to the intensity data of this compound using the SQUEEZE program of the Platon software suite.²⁸ The Flack parameter for [PtCl(PMe₃)₃][BCl₄] indicates that this crystal may be a racemic twin. Crystallographic data for all compounds are presented in Table 1.

CCDC reference numbers 618539-618552.

For crystallographic data in CIF or other electronic format see DOI: 10.1039/b612039n

Acknowledgements

We thank the EPSRC for support and the ORS Scheme for a studentship (CF).

References

- (a) S. Aldridge and D. L. Coombs, Coord. Chem. Rev., 2004, 248, 535; (b) H. Braunschweig and M. Colling, Coord. Chem. Rev., 2001, 223, 1; (c) H. Braunschweig, Angew. Chem., Int. Ed., 1998, 37, 1786; (d) G. J. Irvine, M. J. G. Lesley, T. B. Marder, N. C. Norman, C. R. Rice, E. G. Robins, W. R. Roper, G. R. Whittell and L. J. Wright, Chem. Rev., 1998, 98, 2685; (e) H. Wadepohl, Angew. Chem., Int. Ed. Engl., 1997, 36, 2441; (f) M. R. Smith III, Prog. Inorg. Chem., 1999, 48, 505.
- 2 For recent reviews on diboration, see: (a) T. B. Marder and N. C. Norman, *Top. Catal.*, 1998, **5**, 63; (b) T. Ishiyama and N. Miyaura, J. Synth. Org. Chem. Jpn., 1999, **57**, 503; (c) T. Ishiyama and N. Miyaura, J. Organomet. Chem., 2000, **611**, 392; (d) T. Ishiyama and N. Miyaura, *Chem. Rec.*, 2004, **3**, 271. For recent references on C-H borylation see, for example:; T. Ishiyama and N. Miyaura, J. Organomet. Chem., 2003, **680**, 3; D. N. Coventry, A. S. Batsanov, A. E. Goeta, J. A. K. Howard, T. B. Marder and R. N. Perutz, Chem. Commun., 2005, 2172 and references therein.
- 3 See: G. Schmid, Angew. Chem., Int. Ed. Engl., 1970, 9, 819 and references therein.
- 4 G. Schmid, W. Petz, W. Arloth and H. Nöth, *Angew. Chem., Int. Ed. Engl.*, 1967, **6**, 696.
- 5 G. Scmid, Chem. Ber., 1969, 102, 191.
- 6 A. Kerr, T. B. Marder, N. C. Norman, A. G. Orpen, M. J. Quayle, C. R. Rice, P. L. Timms and G. R. Whittell, *Chem. Commun.*, 1998, 319; N. Lu, N. C. Norman, A. G. Orpen, M. J. Quayle, P. L. Timms and G. R. Whittell, *J. Chem. Soc., Dalton Trans.*, 2000, 4032.
- 7 T. H. Peterson, J. T. Golden and R. G. Bergman, *Organometallics*, 1999, **18**, 2005.
- 8 G. J. Irvine, C. E. F. Rickard, W. R. Roper, A. Williamson and L. J. Wright, Angew. Chem., Int. Ed., 2000, 39, 948.
- 9 C. E. F. Rickard, W. R. Roper, A. Williamson and L. J. Wright, *Organometallics*, 2002, 21, 1714; C. E. F. Rickard, W. R. Roper, A. Williamson and L. J. Wright, *Organometallics*, 2002, 21, 4862; G. R. Clark, G. J. Irvine, W. R. Roper and L. J. Wright, *J. Organomet. Chem.*, 2003, 680, 81.
- 10 S. Aldridge, R. J. Calder, R. E. Baghurst, M. E. Light and M. B. Hursthouse, J. Organomet. Chem., 2002, 649, 9.
- 11 A. A. Dickinson, D. J. Willock, R. J. Calder and S. Aldridge, Organometallics, 2002, 21, 1146.
- 12 H. Braunschweig, K. Radacki, F. Seeler and G. R. Whittell, *Organometallics*, 2004, 23, 4178.
- 13 H. Braunschweig, K. Radacki, D. Rais and G. R. Whittell, Angew. Chem., Int. Ed., 2005, 44, 1192.
- 14 H. Braunschweig, K. Radacki, D. Rais, F. Seeler and K. Uttinger, J. Am. Chem. Soc., 2005, 127, 1386.
- 15 H. Braunschweig, K. Radacki, D. Scheschkewitz and G. R. Whittell, Angew. Chem., Int. Ed., 2005, 44, 1658.
- 16 W. Clegg, F. J. Lawlor, G. Lesley, T. B. Marder, N. C. Norman, A. G. Orpen, M. J. Quayle, C. R. Rice, A. J. Scott and F. E. S. Souza, J. Organomet. Chem., 1998, 550, 183.
- 17 D. Curtis, M. J. G. Lesley, N. C. Norman, A. G. Orpen and J. Starbuck, J. Chem. Soc., Dalton Trans., 1999, 1687.
- 18 H. Braunschweig, K. Radacki, D. Rais and F. Seeler, Organometallics, 2004, 23, 5545.
- 19 T. R. Durkin and E. P. Schram, *Inorg. Chem.*, 1972, **11**, 1054.
- 20 M. Fishwick, H. Nöth, W. Petz and M. G. H. Wallbridge, *Inorg. Chem.*, 1976. **15**, 490.
- 21 J. Zhu, Z. Lin and T. B. Marder, Inorg. Chem., 2005, 44, 9384.

22 G. Lesley, P. Nguyen, N. J. Taylor, T. B. Marder, A. J. Scott, W. Clegg and N. C. Norman, *Organometallics*, 1996, 15, 5137.
L. E. Crascall and J. L. Spencer, *Inorg. Synth.*, 1990, 28, 127.

 25 L. E. Crascall and J. L. Spencer, *Inorg. Synth.*, 1990, 26, 127.
24 C. N. Welch and S. G. Shore, *Inorg. Chem.*, 1968, 7, 225; R. J. Brotherton and W. G. Woods, *US Patent*, 3009 941, 1961; F. J. Lawlor, N. C. Norman, N. L. Pickett, E. G. Robins, P. Nguyen, G. Lesley, T. B. Norman, N. L. Pickett, E. G. Robins, P. Nguyen, G. Lesley, T. B. Norman, N. L. Pickett, E. G. Robins, P. Nguyen, G. Lesley, T. B. Norman, N. L. Pickett, E. G. Robins, P. Nguyen, G. Lesley, T. B. Norman, N. L. Pickett, E. G. Robins, P. Nguyen, G. Lesley, T. B. Norman, N. L. Pickett, E. G. Robins, P. Nguyen, G. Lesley, T. B. Norman, N. L. Pickett, E. G. Robins, P. Nguyen, G. Lesley, T. B. Norman, N. L. Pickett, E. G. Robins, P. Nguyen, G. Lesley, T. B. Norman, N. L. Pickett, E. G. Robins, P. Nguyen, G. Lesley, T. B. Norman, N. L. Pickett, E. G. Robins, P. Nguyen, G. Lesley, T. B. Norman, N. L. Pickett, E. G. Robins, P. Nguyen, G. Lesley, T. B. Norman, N. L. Pickett, E. G. Robins, P. Nguyen, G. Lesley, T. B. Nguyen, G. Lesley, G. Lesley, G. Lesley, G. Lesley, T. B. Nguyen, G. Lesley, G. Lesley, G. Lesley, G. Marder, J. A. Ashmore and J. C. Green, *Inorg. Chem.*, 1998, **37**, 5282; W. Clegg, M. R. J. Elsegood, F. J. Lawlor, N. C. Norman, N. L. Pickett, E. G. Robins, A. J. Scott, P. Nguyen, N. J. Taylor and T. B. Marder,

Inorg. Chem., 1998, 37, 5289; M. J. G. Lesley, N. C. Norman and C. R. Rice, Inorg. Synth., 2004, 34, 1.

- 25 SAINT integration software, Bruker-AXS Inc., Madison, WI, 2004.
- 26 G. M. Sheldrick, SADABS absorption correction program, University of Göttingen, Germany, 2003.
- 27 SHELXTL program system, Bruker-AXS Inc., Madison, WI, 2003.
- 28 A. L. Spek. PLATON, A Multipurpose Crystallographic Tool, Utrecht University, Utrecht, The Netherlands, 2005.